

Absolute dimensions of the unevolved B-type eclipsing binary GG Orionis¹

Guillermo Torres

Harvard-Smithsonian Center for Astrophysics, 60 Garden St., Cambridge, MA 02138

Claud H. Sandberg Lacy²

Department of Physics, University of Arkansas, Fayetteville, AR 72701

Antonio Claret

Instituto de Astrofísica de Andalucía, CSIC, Apartado 3004, E-18080 Granada, Spain

Jeffrey A. Sabby²

Department of Physics, University of Arkansas, Fayetteville, AR 72701

To appear in *The Astronomical Journal*, December 2000

¹Some of the observations reported here were obtained with the Multiple Mirror Telescope, a joint facility of the Smithsonian Institution and the University of Arizona.

²Visiting astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatories, operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation.

ABSTRACT

We present photometric observations in B and V as well as spectroscopic observations of the detached, eccentric 6.6-day double-lined eclipsing binary GG Ori, a member of the Orion OB1 association. Absolute dimensions of the components, which are virtually identical, are determined to high accuracy (better than 1% in the masses and better than 2% in the radii) for the purpose of testing various aspects of theoretical modeling. We obtain $M_A = 2.342 \pm 0.016 M_\odot$ and $R_A = 1.852 \pm 0.025 R_\odot$ for the primary, and $M_B = 2.338 \pm 0.017 M_\odot$ and $R_B = 1.830 \pm 0.025 R_\odot$ for the secondary. The effective temperature of both stars is 9950 ± 200 K, corresponding to a spectral type of B9.5. GG Ori is very close to the ZAMS, and comparison with current stellar evolution models gives ages of 65-82 Myr or 7.7 Myr depending on whether the system is considered to be burning hydrogen on the main sequence or still in the final stages of pre-main sequence contraction. Good agreement is found in both scenarios for a composition close to solar. We have detected apsidal motion in the binary at a rate of $\dot{\omega} = 0.00061 \pm 0.00025$ deg cycle $^{-1}$, corresponding to an apsidal period of $U = 10700 \pm 4500$ yr. A substantial fraction of this ($\sim 70\%$) is due to the contribution from General Relativity, and our measurement is entirely consistent with theory. The eccentric orbit of GG Ori is well explained by tidal evolution models, but both theory and our measurements of the rotational velocity of the components are as yet inconclusive as to whether the stars are synchronized with the orbital motion.

Subject headings: binaries: eclipsing — binaries: spectroscopic — stars: evolution — stars: fundamental parameters — stars: individual (GG Orionis)

1. Introduction

The discovery of GG Orionis (HD 290842, Tycho 4767 857 1, $V = 10.4 - 11.1$, B9.5 V, $\alpha = 05^h43^m10^s.2$, $\delta = -00^\circ41'15''$, epoch and equinox J2000) as a variable star is due to Hoffmeister (1934), who observed the object photographically at the Sonneberg Observatory. The correct period of 6.631 days was first given by Kordylewski (1951) based on visual and photographic times of minimum. This author obtained a mean visual light curve, and established that the orbit is eccentric from the displacement of the secondary minimum.

Aside from the occasional measurement of the times of eclipse by a number of authors, GG Ori has remained until recently a rather neglected system. Double lines in the spectrum were detected by Lacy (1984), who described them as being narrow and of nearly equal strength, but no detailed spectroscopic study has been made to date. The first photoelectric light curves were published by Zakirov (1997), who presented light elements for this well detached binary solved by the method of Lavrov (1993).

Based on the fact that the orbit is eccentric, it is expected that the system may present a measurable apsidal motion. This effect is of great interest in the study of detached eclipsing binaries because it provides information on the internal structure of stars that may be compared with predictions from theory. From its spectral type and other known properties, GG Ori was listed by Giménez (1985) as a good candidate for the study of the contribution of General Relativity to the secular displacement of the line of apsides, given that the relativistic effect is expected to be dominant in this particular case.

GG Ori is located in the Orion OB1 association (see, e.g., Blaauw 1964; Warren & Hesser 1977), a complex region of star formation that has been the subject of numerous studies to determine the properties of the population of young stars and surrounding gas. The binary is located not far from the Belt of Orion, and therefore there is reason to expect that the system might also be quite young, adding to its interest.

In this paper we present new high-quality photoelectric light curves in two passbands, which we analyze together with other published photometry. We also report the results of our intensive spectroscopic monitoring of GG Ori that, combined with the light curves, enable us to derive highly precise absolute dimensions for both components of the system. The stars turn out to be nearly identical in all their properties.

Our current knowledge of the internal structure and evolution of stars is such that observed stellar properties determined with errors of even 5% are of little value to constrain the models in any significant way, since the differences between competing calculations or the effects of slight changes in the input physics or atomic constants are below this level

(see Andersen 1991, 1998). The masses we obtain here have errors smaller than 1%, and the radii are determined to better than 2%. We use these determinations for a comparison with the predictions of recent stellar evolution models, which suggest a very young age for the system. An analysis of all available eclipse timings along with our radial velocities leads to a small but apparently significant apsidal motion detection. We discuss the importance of the relativistic effect mentioned above compared to classical effects from rotational and tidal distortions. In addition, we examine the predictions from tidal theory regarding spin-orbit synchronization and the circularization of the orbit.

2. Spectroscopic observations and reductions

GG Ori was observed spectroscopically at the Kitt Peak National Observatory (KPNO) and at the Harvard-Smithsonian Center for Astrophysics (CfA). Observations at KPNO were made with the 2.1m and coudé-feed spectrometer from 1983 December to 1999 March. A total of 15 spectrograms were obtained with a variety of CCD detectors. A spectral resolution of 0.02 nm or 2 pixels was used before May 1998, and 0.03 nm or 3 pixels after that date. The spectral coverage was about 10 nm in the first 12 spectrograms, and 32 nm in the last 3 spectrograms. The central wavelength of those observations was 450 nm.

Projected rotational velocities ($v \sin i$) were determined by comparing line widths of the binary components’ unblended features with artificially-broadened features in the spectra of comparison stars with known values of $v \sin i$. These comparison stars were chosen to be of nearly the same spectral type as GG Ori, and were observed with the same instrumental configuration as the binary. The reference stars used were *o* Peg (HR 8641, A1 IV, $v \sin i = 10 \text{ km s}^{-1}$; Fekel 1998, private communication), 68 Tau (HR 1389, A2 IV-V, $v \sin i = 18 \text{ km s}^{-1}$; Hoffleit 1982), and HR 8404 (21 Peg, B9.5 V, $v \sin i = 4 \text{ km s}^{-1}$; Fekel 1999). From these comparisons in 10 of the spectrograms, values of $24 \pm 2 \text{ km s}^{-1}$ and $23 \pm 2 \text{ km s}^{-1}$ were determined for the primary and secondary in GG Ori, respectively, based mainly on the Mg II 448.1 nm line. The uncertainties given account for the scatter of the line width measurements as well as the agreement between the results using different standard stars. The component of GG Ori we call here the “primary” (also star “A”) is the one eclipsed at phase 0.0 in the light curve (see §4). Formally it is also the more massive one, but only marginally so, as we describe later.

Radial velocities of the components were measured by cross-correlation with the FXCOR task in IRAF³, using standard stars (same as above) with known radial velocities

³IRAF is distributed by the National Optical Astronomy Observatories, which is operated by the

as templates. Radial velocities for the low-amplitude spectroscopic binary star *o* Peg were taken from the orbit of Fekel (1999). The radial velocity of HR 8404 was taken as $+0.2 \text{ km s}^{-1}$ based on 12 velocities by Fekel (1999), and for 68 Tau we adopted the value $+39.0 \text{ km s}^{-1}$, from the same source. Our measurements for GG Ori from the 11 spectra with relatively unblended lines are given in Table 1.

In addition to the radial and rotational velocities, we determined the line strength ratio from 17 line pairs in spectra of good signal-to-noise (S/N) ratio. This may be used as a proxy for the light ratio between the stars. We obtained $\text{EW}_A/\text{EW}_B = 1.01 \pm 0.02$, corresponding to the blue region of the spectrum. Because the two components are virtually identical, no correction for the difference in temperature is needed, and the ratio of the equivalent widths, EW_A/EW_B , can be assumed to be identical to the light ratio.

Our observations of GG Ori at the CfA were collected mostly with the echelle spectrograph on the 1.5m Tillinghast reflector at the F. L. Whipple Observatory on Mt. Hopkins (Arizona), where the system was monitored from 1996 March to 2000 March. Occasionally we observed with an identical spectrograph on the Multiple Mirror Telescope (also atop Mt. Hopkins, Arizona). A total of 42 spectra were recorded with a photon-counting Reticon detector covering a single echelle order centered at 518.7 nm. The resolution is 0.015 nm ($\lambda/\Delta\lambda \sim 35,000$), and the spectra span about 4.5 nm. The S/N ratios range from about 25 to 40 per resolution element. The zero point of the velocity system was monitored by means of nightly exposures of the dusk and dawn sky, in the manner described by Latham (1992). The accuracy of the CfA velocity system, which is within about 0.1 km s^{-1} of the reference frame defined by minor planets in the solar system, is documented in the previous citation and also by Stefanik et al. (1999).

Radial velocities were determined using the CfA implementation of TODCOR (Zucker & Mazeh 1994), a two-dimensional cross-correlation algorithm that uses two templates, one for each component of the binary. The templates were selected from a large library of synthetic spectra based on the latest model atmospheres by R. L. Kurucz, computed for us specifically for the wavelength of our observations (Morse & Kurucz 1998, private communication). The instrumental profile is explicitly included by applying a Gaussian convolution with a FWHM of 8.5 km s^{-1} , corresponding to our spectral resolution. These synthetic spectra have been calculated over a wide range of effective temperatures, projected rotational velocities, metallicities, and surface gravities. Initially we used templates with a temperature of $T_{\text{eff}} = 10000 \text{ K}$ for both stars, surface gravity of $\log g = 4.5$ (cgs), and

solar metallicity. The projected rotational velocities of the components were determined by running extensive grids of correlations against templates covering a range of values of $v \sin i$, seeking the best match to the observed spectra as determined by the correlation coefficient averaged over all exposures. We obtained $v \sin i = 16 \pm 1 \text{ km s}^{-1}$ for both components, which is significantly different than our determination based on KPNO spectra. We discuss this further in §5 and §6. For the final templates we adopted a temperature of 9950 K for both stars and $\log g = 4.3$, based on the results from the light curve solutions and spectroscopic orbits presented below.

Systematic errors in the radial velocities resulting from line blending are always a concern, particularly when the goal is to achieve the highest possible precision and accuracy in the mass determinations. Although the use of a two-dimensional correlation technique such as TODCOR in principle minimizes those errors (see Latham et al. 1996), experience has shown us that this must be checked on a case-by-case basis, particularly in view of the narrow spectral window of the CfA observations (e.g., Torres et al. 1997; 2000). For this we generated artificial composite spectra by adding together synthetic spectra for the two components with Doppler shifts appropriate for each actual time of observation, computed from a preliminary orbital solution. We adopted also a light ratio close to that for the real stars (see below). We then processed these simulated spectra with TODCOR in the same manner as the real spectra, and compared the input and output velocities. The differences derived in this way vary systematically with phase (or radial velocity), as expected, and were applied to the real velocities as corrections, even though in the case of GG Ori they are quite small ($\leq 0.5 \text{ km s}^{-1}$) and hardly affect the results. The final radial velocities from the CfA spectra with the corrections included are given in Table 2.

The light ratio was derived from these observations using TODCOR as described by Zucker & Mazeh (1994). Because of the relatively small number of lines in the 4.5 nm spectral window observed at CfA, significant errors can be introduced in the light ratio when features of each component shift in and out of this region by up to 0.2 nm due to orbital motion. To estimate and correct for this effect we followed a procedure analogous to that described above for the velocities, and found the magnitude of the systematic error to be about 3.5%. After correcting for this we obtain $L_A/L_B = 1.05 \pm 0.03$ at a mean wavelength of 518.7 nm, which is sufficiently close to the visual band that we will assume it is the light ratio in V , given that the stars are virtually identical.

Preliminary double-lined orbital solutions based on our radial velocities were computed separately for the KPNO and CfA measurements, and are compared in Table 3. Initially we adopted the eclipse ephemeris given in the General Catalog of Variable Stars (GCVS) (Kholopov 1985): $\text{Min I (HJD)} = 2,433,596.496 + 6.63147 \cdot E$. However, the residuals from

the orbit for the CfA data showed a systematic pattern and the scatter was significantly larger than expected from previous experience with similar spectroscopic material. This suggested that the ephemeris is not accurate enough to be propagated forward by ~ 50 yr (2600 cycles) to the epoch of our velocity measurements. Therefore, a new linear ephemeris computed from times of minimum collected from the literature was derived (see next section) and used here. The possibility of a systematic difference between the center-of-mass velocities of the primary and secondary components was explored by solving for an offset simultaneously with the orbital elements. This was done independently for the two data sets. No shift was expected because the stars are so similar. The results in the sense $\langle \text{primary minus secondary} \rangle$ were $+1.42 \pm 0.97 \text{ km s}^{-1}$ for KPNO and $-0.74 \pm 0.53 \text{ km s}^{-1}$ for CfA. Since these are indeed not significantly different from zero, the velocities for the two components can be considered for all practical purposes to be on the same reference frame.

As seen in Table 3, there are slight differences in some of the elements of these two orbital solutions, in particular in the velocity amplitude of the secondary component (K_B), but we do not consider them to be significant in view of the small number of KPNO observations (only 11). Therefore, for the final solution discussed below we have merged the two data sets.

3. Apsidal motion analysis and final spectroscopic solution

As indicated earlier, the noticeable eccentricity of the orbit of GG Ori suggests the possibility of apsidal motion, even though none has been reported for this system. In extreme examples this can affect the spectroscopic solutions quite significantly, as in the case of V477 Cyg (Popper 1968). Before computing the final orbital fit, we therefore searched the literature for eclipse timings, both to improve the ephemeris and to investigate the apsidal motion.

Table 4 lists all times of minimum available to us, of which there are 25 primary minima and 32 secondary minima covering about 65 yr (1930-1995). The majority were determined by visual or photographic means, and only the 6 recent photoelectric timings and one visual timing have published uncertainties. For the rest we determined the errors iteratively, based on the mean residuals for each type of observation from preliminary fits: $\sigma_{\text{pg}} = 0.038$ days for the photographic minima, and $\sigma_v = 0.028$ days for the visual minima. One visual estimate was found to give a large residual (0.14 days, $\sim 5\sigma$), and was excluded. It is indicated in parentheses in Table 4.

A linear ephemeris fit to these data gives $\text{Min I (HJD)} = 2,449,717.66253(21) +$

$6.6314948(16) \cdot E$, with a phase difference between the primary and secondary minima of $\Delta\Phi = 0.42252 \pm 0.00024$. This is the ephemeris we used above in our preliminary orbital solutions, and the period is very similar to that reported by Zakirov (1997) based on a smaller number of observations. Although no unusual trends were seen in the timing residuals, we investigated the possibility of apsidal motion by computing the apparent periods separately for the primary and the secondary. We obtained $P_I = 6.6314971 \pm 0.0000021$ days and $P_{II} = 6.6314890 \pm 0.0000030$ days, which are different at the 2.2σ level. Next we submitted these same data to an apsidal motion analysis using the method by Lacy (1992b). For this we assumed a fixed eccentricity equal to the average of the values derived in columns (2) and (3) of Table 3, and a fixed inclination angle of 89° from preliminary light curve solutions. The result is only marginally significant ($\dot{\omega} = 0.00050 \pm 0.00027$ deg cycle $^{-1}$), largely due to the weak constraint provided by the early photographic and visual times of minimum, which have relatively low weight. Changes in those weights have little effect.

Our radial velocities extend the time base provided by the times of minimum by another 5 yr, and contain valuable information on the longitude of periastron that can also be used. In addition, they provide a much stronger constraint on the eccentricity, which must otherwise be fixed when using only eclipse timings. The optimal solution, therefore, is to combine both kinds of measurements into a single least squares fit, solving simultaneously for the spectroscopic orbital elements and the apsidal motion. In this way the information on the period contained in the times of minimum is implicitly taken into account for the spectroscopic elements, rather than having to adopt a fixed ephemeris as we have done in §2. At the same time, the effect of the possible rotation of the line of apsides on the spectroscopic elements is also accounted for. In addition, we have allowed for an arbitrary shift between the KPNO and CfA velocity systems in view of the difference in the way the two zero points were established. The offset turns out to be negligible: $\langle \text{CfA-KPNO} \rangle = -0.06 \pm 0.39$ km s $^{-1}$.

The results of this simultaneous fit are given in column (4) of Table 3. The relative weights of the primary and secondary velocities in each data set, as well as the weights of the times of minimum, have been iterated until reaching convergence for a reduced χ^2 of unity. Tests show once again that the weights assigned to the visual and photographic timings, or to the velocities, have little effect on the results. The velocity residuals from this solution are listed separately in Table 1 and Table 2, and the timing residuals are given in Table 4. The final spectroscopic orbital solution is represented graphically in Fig. 1, along with the observations.

The apsidal motion resulting from the final fit is $\dot{\omega} = 0.00061 \pm 0.00025$ deg cycle $^{-1}$, which is significant at the 2.5σ level, and differs from our preliminary estimate above by

less than half of its uncertainty. Also listed in Table 4 are the sidereal and anomalistic periods (P_s and P_a), as well as the apsidal motion period $U = 10700 \pm 4500$ yr. Periastron passage occurs at a photometric phase of 0.06. A plot of the $O-C$ deviations of the times of minimum from the linear terms of the apsidal motion is given in Fig. 2 along with the predicted deviations. The top panel draws attention to the fact that the observations cover less than 1% of a full cycle, while the bottom panel expands the region around the observations, with the coverage provided by the radial velocity measurements being indicated in the upper right.

4. Photometric observations and light curve solutions

Differential light curves were obtained by CHSL at the Cerro Tololo Inter-American Observatory (CTIO) with the 24-inch Lowell telescope in the Johnson B and V bands during 1993-1995. Absolute indices were also measured separately based on measurements that were carefully tied to the standard UBV system through observations of secondary standards from Landolt (1973), typically 30-40 standards per night. The absolute photometry was made at Mount Laguna Observatory near San Diego (California) in the fall season of 1989, and at CTIO in 1988-1990 and 1993-1995. The procedures that were used are described in detail by Lacy (1992a). The UBV indices on the standard system are shown in Table 5, and the 257 differential BV observations are given in Table 6 and Table 7. The differential magnitudes are in the sense $\langle \text{variable} - \text{comparison} \rangle$ and are referred to BD $-1^\circ 1013$ (HD 38165, B9 V). The comparison stars (see Table 5) were found to be constant at a level of about 0.008 mag for the standard deviation of the differences between comparison stars. The precision of the CTIO differential observations, based on previous results, is estimated to be about 0.006 mag in both B and V at the magnitudes of the program stars.

In addition to the absolute photometric indices mentioned above, which we have computed separately for the two observing intervals, $uvby$ indices are available from Hilditch & Hill (1975). The results from the three different sources are somewhat inconsistent. The V magnitude is slightly brighter by about 2.5% in the 1988-1990 results compared to the 1993-1995 results and the value of Hilditch & Hill (1975), which is $V = 10.380 \pm 0.015$. The $B-V$ color index during 1993-1995 is considerably bluer than the value measured in 1988-1990. These discrepancies cannot be explained by differences in sky conditions (which were always photometric), nor by differences in aperture size, centering errors, or seasonal variations. Furthermore, the scatter from the light curve fits described later in this section is significantly larger than expected based on our previous results with the equipment used

at CTIO at these magnitude levels. The standard deviation of the residuals from the fitted photometric orbit of FS Mon (Lacy et al. 2000), for example, was 0.006 and 0.005 mag in B and V , respectively, whereas the corresponding figures for GG Ori (see below for details) are 0.014 and 0.009 mag. Both binaries were observed during the same runs. Intrinsic variability of one or both of the components of GG Ori is therefore suspected.

Differential light curves in the Johnson U , B , V , and R bands have also been obtained in Uzbekistan by the group led by M. Zakirov. They used 0.6m telescopes at the Maidanak Observatory, in the southern part of Uzbekistan. Their differential magnitudes in the sense ⟨variable–comparison⟩ relative to the star BD $-1^{\circ}1013$ have been described by Zakirov (1997). The transformation of these measurements to the standard system is somewhat less secure than for our own observations, but there are marginal indications that the $B-V$ index of GG Ori became bluer by a few hundredths of a magnitude in the interval 1993-1994, toward the end of their observations. Though far from being conclusive evidence of intrinsic variability, we note that this is in the same direction as the trend suggested in Table 5. We have re-analyzed the differential observations by Zakirov with the same methods we use for our own measurements.

The light curve fitting was done with the NDE model as implemented in the code EBOP (Etzel 1981; Popper & Etzel 1981), and the ephemeris adopted is that of §3. The main adjustable parameters are the ratio of the central surface brightness of the secondary star (J_B) in units of that of the primary, the relative radius of the primary (r_A) in units of the separation, the ratio of the radii ($k \equiv r_B/r_A$), the inclination of the orbit (i), and the geometric factors $e \cos \omega$ and $e \sin \omega$ which account for the orbital eccentricity. As usual, we have allowed also for a photometric scale factor and a phase shift in all solutions. Auxiliary quantities needed in the analysis include the gravity-brightening coefficients, for which we adopt the values 1.00, 0.85, 0.70, and 0.59 in U , B , V , and R , respectively, from Martynov (1973). For the reflection coefficients we adopted the value 1.0, as appropriate for stars with radiative envelopes. Limb-darkening coefficients (u) were included as one of the variables to be fitted, although in one case (noted below) it was fixed to an appropriate value from the tables of Wade and Rucinski (1985). The mass ratio ($q \equiv M_B/M_A = 0.9982$) was adopted from the spectroscopic analysis in §3.

Preliminary solutions revealed a few outliers with residuals greater than 5σ (one of the CTIO measurements in B , two in V , and one of Zakirov’s measurements in each of his four bands), which were given zero weight in subsequent iterations.

The fitting procedure converged in the general solutions of the CTIO light curves with all variables adjusted simultaneously (see columns 2 and 3 in Table 8). As is often the case in the analysis of partially eclipsing binary stars with nearly equal components, the fitted

values of the parameter k , the ratio of radii, are rather uncertain, but near unity. Attempts at general solutions of the Zakirov light curves failed to converge. In cases such as this light ratios estimated from spectrograms are very important for constraining the photometric solutions. For GG Ori we have at our disposal two spectroscopic determinations of the light ratio, one approximately in the B band (from KPNO) and the other close to the V band (from CfA). A grid of EBOP solutions was run on the CTIO light curves for fixed values of k between 0.94 and 1.10, in steps of 0.01. Within this range of about 15%, the rms residuals from the fits were found to change very little as a function of k (≤ 0.00005 mag in B and ≤ 0.00006 mag in V), supporting our concerns about the indeterminacy of k from the light curves alone. The sum of the relative radii changes by less than 0.7% over the entire range of k values. The light ratios in B and V were computed from each of these solutions, and then by interpolation we determined the k values that correspond to our measured spectroscopic light ratios: $k = 0.996 \pm 0.011$ from the KPNO spectra, and $k = 0.972 \pm 0.015$ from the CfA spectra (see Fig. 3). The value we adopt is the weighted average, $\langle k \rangle = 0.988 \pm 0.009$. For comparison, the average ratio of the radii derived from the CTIO light curve solutions without the constraint from spectroscopy is $k = 1.03 \pm 0.04$ (see Table 8), which is different by 1σ .

With the ratio of the star sizes fixed to the weighted average given above, we repeated the light curve fits separately for the CTIO and Zakirov (1997) photometry, and the adopted solutions are given in Table 8 and Table 9. Attempts to include third light as a parameter showed that, within uncertainties of about 1%, it was not significantly different from zero.

The CTIO observations in B and V and the best fit models are shown in Fig. 4, and the data by Zakirov (1997) in $UBVR$ along with the corresponding fitted light curves are shown in Fig. 5. Table 10 gives the weighted averages of the light elements in each band obtained from the two data sets. In addition to r_A and k , we list also the sum of the relative radii, $r = r_A + r_B$, and the radius of the secondary, r_B . The errors for these quantities were derived as described in the Appendix. There is good agreement between the photometric determinations of $e \cos \omega$ and $e \sin \omega$ and the corresponding spectroscopic values, which are -0.1200 ± 0.0017 and $+0.1865 \pm 0.0020$, respectively.

The results in Table 10 show that the components of GG Ori are nearly indistinguishable in size and of virtually the same luminosity, which is consistent with the fact that they are also very similar in mass (§3). The departure from the spherical shape is insignificant. Both eclipses are partial, with approximately 91% of the light of the primary blocked at phase 0.0 (which is a transit), and 90% of the light of the secondary eclipsed at the other minimum. The earlier study by Zakirov (1997) reported the primary eclipse to be total (larger secondary star blocking the primary), although the ratio of the radii was much

closer to 1.0.

The uncertainties assigned to the adjusted quantities in Table 10 are generally the internal errors produced by EBOP. As is well known, the formal uncertainties in least-squares solutions in which one or more quantities have been held fixed are typically too small because correlations between the elements are artificially eliminated (as we have done by fixing k , for example). The quantities k , r_A , and i are the most important for determining the absolute dimensions of the components. The error in k is based on the uncertainty in the spectroscopic light ratios, and therefore does not suffer from this shortcoming. For r_A and i we have increased the formal uncertainties to account for all contributions to the error.

The rms deviations given in Table 8 for the CTIO observations are somewhat larger than expected, based on previous experience, and suggest that one or both components of GG Ori may be intrinsically variable. We have searched for patterns in the residuals by computing the power spectra in the B and V bands, but no significant periodicities were found.

5. Absolute dimensions

The combination of the spectroscopic results in Table 3 and the light curve results in Table 10 leads to the absolute masses and radii for GG Ori, shown in Table 11. The masses are determined with a precision of 0.7%, and the radii are good to about 1.4%.

The fact that the surface brightness ratio J_B is near unity over a wide range of wavelengths indicates that the effective temperatures of the stars are essentially the same. From J_B in the V band the difference in visual surface brightness is $\Delta F'_V = 0.0002 \pm 0.0004$ (Popper 1980), which translates into a completely negligible color difference $\Delta(B-V) = 0.0005 \pm 0.0009$, and a formal temperature difference of only 8 K.

The discrepancies in the absolute photometry at different epochs pointed out earlier in §4 complicate the derivation of the effective temperature of the components. Because the light variations may be real, we have chosen to adopt the straight average of the largest and smallest UBV indices for GG Ori in Table 5: $V = 10.372 \pm 0.013$, $B - V = 0.511 \pm 0.021$, $U - B = 0.324 \pm 0.004$. The errors for the first two of these quantities are simply half of the difference between the 1988-1990 and 1993-1995 measurements. The reddening was derived using the reddening-free index $Q = (U - B) - 0.72(B - V)$ (Johnson & Morgan 1953) and the calibration by Deutschman, Davis & Schild (1976). We obtained $(B - V)_0 = -0.036 \pm 0.006$, $(U - B)_0 = -0.070 \pm 0.020$, $E(B - V) = 0.547 \pm 0.022$, and $A_V = 1.696 \pm 0.068$ (using

$A_V = 3.1 \cdot E(B-V)$). The formal errors do not account for uncertainties in the calibrations themselves, which are difficult to quantify. These color indices correspond to the combined light, but since the components are virtually identical they are also the indices of the individual stars.

An independent measure of the reddening may be obtained from the *uvby* photometry by Hilditch & Hill (1975): $b-y = 0.408 \pm 0.015$, $c_1 = 1.002 \pm 0.030$, $m_1 = +0.016 \pm 0.020$. The calibration by Crawford (1978) for B-type stars leads to $(b-y)_0 = -0.030$ and $E(b-y) = 0.438$, from which $E(B-V) = 0.592$. This is slightly larger than the reddening derived from the *UBV* photometry. However, GG Ori is near the end of the range of validity of the *uvby* calibrations, where they become rather uncertain. Also, we note that the $(b-y)_0$ index corresponds to a spectral type of B8.5 according to Table 1 by Popper (1980). But the fact that we see no sign of the He I 447.1 nm line in our KPNO spectra of the object strongly suggests that the spectral type cannot be earlier than B9.5. We have therefore chosen to rely only on the Johnson photometry above.

The de-reddened $B-V$ index corresponds to a temperature of 9950 ± 200 K, and a spectral type of B9.5 (Popper 1980). Further properties of the stars are listed in Table 11, including the mean density ($\bar{\rho}$), the absolute visual magnitudes, and the distance. The latter two are based on the visual surface brightness parameter $F'_V = 3.974 \pm 0.003$ derived from the intrinsic color $(B-V)_0$ and the tabulation by Popper (1980), and are thus independent of bolometric corrections.

The distance modulus of GG Ori, $m-M = 8.20 \pm 0.10$ ($d = 438 \pm 20$ pc), is similar to other determinations for various subregions of the Orion OB1 complex (Warren & Hesser 1977; Brown, de Geus, & de Zeeuw 1994; Brown, Walter & Blaauw 1999; de Zeeuw et al. 1999), and lends support to its association.

Also listed in Table 11 are the projected rotational velocities expected if the axial rotations were synchronized with the mean orbital motion of the binary and with the orbital motion at periastron (pseudo-synchronization). The two independent measurements of $v \sin i$ in §2 based on our KPNO and CfA spectra disagree, with the KPNO results being closer to the pseudo-synchronous values, while the CfA determinations suggest synchronization with the mean orbital motion. We discuss this further in §6 in the context of tidal evolution theory.

No metallicity determination is available for GG Ori. A rough estimate can be obtained from our CfA spectra following a procedure analogous to that used to derive $v \sin i$, and the result is $[m/H] = -0.15 \pm 0.20$, consistent with the solar abundance.

6. Discussion

In this section we compare the properties of GG Ori as listed in Table 11 with predictions from models. The three aspects of theory we focus on are stellar evolution, internal structure, and tidal evolution. For each of these we require calculations of the basic properties of the components, which we have taken from evolutionary tracks computed specifically for the exact masses we determined for the stars. The evolutionary code we used is that by Claret (1995). Further details on the input physics are described by Claret & Giménez (1992). Convection in these models is treated with the standard mixing-length prescription, with a fixed mixing-length parameter of $1.52H_p$ that gives the best fit between a solar model and the observed properties of the Sun. A moderate amount of core overshooting is assumed ($\alpha_{ov} = 0.20H_p$), although in the case of GG Ori this effect is insignificant due to the unevolved nature of the system.

6.1. Stellar evolution

Consistent with the young environment surrounding the object (the Orion OB1 association), we find that GG Ori is indeed very close to the Zero Age Main Sequence (ZAMS). This is illustrated in Fig. 6, which shows a greatly enlarged section of the evolutionary tracks as they approach the main sequence. The full extent of the main sequence band is shown in the inset. The binary is so close to the ZAMS that, given the uncertainties, it is difficult to determine whether the stars are already burning hydrogen in their cores, or whether they are still in the final stage of contraction towards the main sequence. The error bars shown reflect the uncertainties in the effective temperatures of the components and their surface gravities. The error in the placement of the tracks due to the uncertainty in the measured masses is indicated in the lower left corner.

The models plotted in Fig. 6 correspond to the case where the system is already on the main sequence (MS). Tracks for two different chemical compositions (Z , Y) are shown, giving metallicities very close to solar ($[m/H] = +0.03$, $dY/dZ = 1.65$, solid lines; and $[m/H] = +0.05$, $dY/dZ = 1.95$, dashed lines) consistent with the indication in §5. The enrichment laws (dY/dZ) implied by the best fit values of the helium abundance (Y) are within the range found in other determinations (see, e.g., Peimbert 1993; Renzini 1994; Pagel & Portinari 1998; Izotov & Thuan 1998), and are also similar to the values favored by other eclipsing binaries (Ribas et al. 2000), using the same set of evolutionary models.

Our attempts to find evolutionary tracks that fit the location of GG Ori assuming it is nearing the end of the pre-main sequence (PMS) phase also gave acceptable results

(see Fig. 7), considering the error bars, although the agreement is formally not as good as the main-sequence case. The PMS tracks shown in the figure are for $[m/H] = +0.07$ and $dY/dZ = 2.09$.

The evolutionary ages derived for the components are quite different in the two scenarios (see Table 12). In the main sequence case the average age for the two stars is 82 Myr (for $Z = 0.020$, $Y = 0.2729$) or 65 Myr (for $Z = 0.021$, $Y = 0.2810$), and the difference in age between the components is similar to the formal uncertainty, which is derived from the error in $\log g$. On the other hand, if GG Ori is on the PMS the models give an average age of only 7.7 Myr. Few cases are known of eclipsing binaries with well-determined absolute dimensions in which at least one component is a bona-fide PMS star: EK Cep (Popper 1987; Claret, Giménez, & Martín 1995), TY CrA (Casey et al. 1998), and possibly RS Cha (Clausen & Nördstrom 1980; Pols et al. 1997; Mamajek, Lawson, & Feigelson 2000). Such systems are particularly valuable to test models of PMS evolution, where theory remains essentially unchallenged by observations so far. The stars in GG Ori are so similar and so close to the ZAMS, however, that the constraint on PMS evolution is not very strong. Unevolved systems such as this are much more useful for testing opacity and metallicity effects in the models when coupled with an accurate spectroscopic determination of the metal abundance, which unfortunately is not yet available for GG Ori. Whether or not the system has already settled on the main sequence, it is undoubtedly young. In this connection we note, incidentally, that light variations such as those hinted at in §4 are not entirely unexpected.

It is a remarkable coincidence that among the eclipsing systems with the best known absolute dimensions, no fewer than *six* have at least one component with virtually the same mass as the stars in GG Ori, to within 1%. These systems are V451 Oph, YZ Cas, β Aur, WX Cep, SZ Cen (Andersen 1991), and V364 Lac (Torres et al. 1999). Thus, a total of eight stars (including GG Ori) can in principle be compared with *a single evolutionary track*, and they span the entire main sequence band from the ZAMS (GG Ori) to the shell hydrogen burning phase (SZ Cen, age ~ 800 Myr). In Fig. 8 we show this comparison against one of the models used above for GG Ori ($Z = 0.020$, $Y = 0.2729$), for a composition close to solar. The agreement with theory is very good, indicating that all these systems are well represented by a single metallicity.

6.2. Internal structure and General Relativity

Our detection of the apsidal motion of the binary in §3 provides the opportunity to test models of the internal structure of stars. In addition, the relatively short orbital period and

large masses of the components along with the fact that the orbit is eccentric led Giménez (1985) to propose GG Ori as a good candidate for the study of the general-relativistic (GR) contribution to the apsidal motion. At the current level of the uncertainties in the measurement of apsidal motion in binaries, the theoretical GR contribution is separable from the classical (Newtonian) terms due to the gravitational quadrupole moment induced by rotation and tides. The total apsidal motion can then be expressed as $\dot{\omega}_{tot} = \dot{\omega}_N + \dot{\omega}_{GR}$. The two effects cannot be measured separately, though, so we have chosen here to compare the total theoretical value with the measured quantity.

In the case of GG Ori, the GR term is predicted to be about 2.5 times larger than the classical term, contributing $\sim 70\%$ to $\dot{\omega}_{tot}$. Following Giménez (1985), we obtain $\dot{\omega}_{GR} = 0.000454 \pm 0.000002 \text{ deg cycle}^{-1}$.

To compute $\dot{\omega}_N$ we have used the internal structure constants from theory (including the higher order terms k_3 and k_4), which are calculated at each point along the evolutionary track of each component. The contribution of the rotational distortions to the Newtonian apsidal motion rate $\dot{\omega}_N$ depends on the ratio between rotational angular velocity and the Keplerian angular velocity. This ratio is often derived under the assumption that the stars are already synchronized at periastron. For a system as young as GG Ori this may not necessarily be the case, and in fact the observational evidence we have is conflicting (§5; see also §6.3). We have therefore relied on our own measurements of the projected rotational velocities, $v \sin i$, even though our two independent determinations from KPNO and CfA spectra are somewhat different (§2). The KPNO values ($v_A \sin i = 24 \pm 2 \text{ km s}^{-1}$, $v_B \sin i = 23 \pm 2 \text{ km s}^{-1}$) lead to $\dot{\omega}_N = 0.000209 \pm 0.000012 \text{ deg cycle}^{-1}$, while the CfA measurements ($v \sin i = 16 \pm 1 \text{ km s}^{-1}$ for both components) give $\dot{\omega}_N = 0.000183 \pm 0.000009 \text{ deg cycle}^{-1}$.

When adding this to the GR term, we obtain $\dot{\omega}_{tot} = 0.000663 \pm 0.000012 \text{ deg cycle}^{-1}$ (KPNO) and $\dot{\omega}_{tot} = 0.000637 \pm 0.000010 \text{ deg cycle}^{-1}$ (CfA), in which the errors account for all contributions from measured quantities. These predictions are to be compared with the observed value, $\dot{\omega}_{obs} = 0.00061 \pm 0.00025 \text{ deg cycle}^{-1}$ (§3). There is good agreement with theory, although the uncertainty in the observed value is large enough that the constraint on the models is very weak. Further measurements of times of minimum are required to improve $\dot{\omega}_{obs}$.

6.3. Tidal evolution

Tidal forces within a binary tend to synchronize the rotation of each component with the orbital motion, and to make the orbit circular. The two main mechanisms proposed

to describe these effects (Tassoul & Tassoul 1997, and references therein; Zahn 1992, and references therein) make somewhat different predictions for the timescales of these processes. In this section we compare our observations with both, although the hydrodynamical mechanism by Tassoul has often been found to be too efficient (see, e.g., Claret, Giménez, & Cunha 1995). Synchronization times and circularization times have been computed as described in the latter reference and also by Claret & Cunha (1997), by integrating the differential equations describing the evolution of the rotation and eccentricity. All time-dependent properties of the stars were interpolated directly from the evolutionary tracks for each component.

In performing these calculations for young stars such as GG Ori a number of complications arise regarding the pre-main sequence phase where the radii of the stars are much larger than on the main sequence. One of them is that in principle the orbital period (or semimajor axis) is also changing during this phase. For example, if the period of GG Ori is assumed to be fixed at the current value of 6.6 days, we find that the size of the orbit is too small to accommodate the very large radii predicted for the stars during the early stages of contraction along the Hayashi tracks. Thus the equations for the orbital evolution (period and eccentricity) and rotational evolution are coupled and must be integrated simultaneously (see, e.g., Duquennoy, Mayor, & Mermilliod 1992).

Zahn & Bouchet (1989) have done this for late type stars ($0.5 M_{\odot} \leq M \leq 1.25 M_{\odot}$) and concluded that all binaries with periods up to 7 or 8 days arrive on the main sequence with their orbits already circularized, and that further evolution of the eccentricity on the main sequence is negligible. However, eccentric systems on the main sequence with periods shorter than this *are* actually observed (see, e.g., Mathieu et al. 1992). For earlier-type binaries such as GG Ori the efficiency of the tidal mechanisms may be different, and a detailed investigation of the evolution in the PMS phase is beyond the scope of this paper.

Tidal theory is still largely in development, and many aspects of it remain somewhat controversial. Nevertheless, useful comparisons with the observations are still possible under certain simplifying assumptions, and have indeed been made in a number of cases. Verbunt & Phinney (1995), Claret, Giménez, & Cunha (1995), Claret & Cunha (1997), and others, have compared the predictions of the main mechanisms mentioned above with the properties of detached eclipsing systems with well-determined absolute dimensions. Their calculations were done in detail (i.e., by *integrating* the corresponding differential equations rather than simply using the *timescales* for synchronization and circularization), but avoided the PMS problem by excluding it altogether (Verbunt & Phinney 1995) or by including only the final stages of contraction, and assumed also that the orbital period does not change significantly, which may be a valid approximation when the stars are already close to the

ZAMS.

We have followed a similar prescription here and considered only the final loop in the evolutionary tracks preceding the ZAMS (see Fig. 8). We start our integrations at the onset of the nuclear reaction $^{12}\text{C}(p, \gamma) ^{13}\text{N}(\beta + \nu) ^{13}\text{C}(p, \gamma) ^{14}\text{N}$. At this stage this is not the only source of energy, since contraction is still taking place. The synchronization and circularization times (t_{sync} , t_{circ}) are not strongly dependent on the chemical composition in the range we have considered above. Using the formalism by Tassoul we obtain $\log t_{sync} = 6.875$ for both components of GG Ori. The predictions from the mechanism by Zahn, on the other hand, give $\log t_{sync}^A = 8.888$ and $\log t_{sync}^B = 8.890$, which are significantly longer. The circularization times are nearly identical in both theories: $t_{circ} = 8.888$ (Tassoul) and $t_{circ} = 8.889$ (Zahn). These values are to be compared with the mean age of the system counted from the starting point of the integrations, which differs by 5.4 Myr from the true ages given in §6.1. The modified age is then $\log t = 7.884$ ($Z = 0.020$, $Y = 0.2729$) or $\log t = 7.775$ ($Z = 0.021$, $Y = 0.2810$), assuming GG Ori is already burning hydrogen on the main sequence.

We illustrate this in a slightly different way in Fig. 9, where we focus on $\log g$ as a sensitive measure of stellar evolution. The value of the surface gravity at which synchronization or circularization is achieved ($\log g_{crit}$) is shown as a function of orbital period for the two main tidal theories. Stars evolve upwards in this diagram. The measured values of $\log g$ for the primary and secondary of GG Ori are represented by crosses, with the error bars being of the same size as the symbols. According to the theory by Zahn the stars should not yet have reached synchronous rotation, which agrees with the measured $v \sin i$ values from our CfA spectra. Also, the orbit should not yet be circular, as is indeed observed. The theory by Tassoul also predicts that the orbit should still be eccentric, but indicates that the stars should already be rotating synchronously with the orbital motion, which appears to be more consistent with the $v \sin i$ values from our KPNO spectra. Although as mentioned above the mechanism by Tassoul is often found to be too efficient, a more accurate measurement of the projected rotational velocities in GG Ori could shed more light on this issue.

7. Conclusions

New photometric and spectroscopic observations of the eccentric binary GG Ori combined with a reanalysis of data from the literature have allowed us to derive definitive orbital parameters and physical properties of the components. Our determinations have formal errors smaller than 1% in the masses and smaller than 2% in the radii. GG Ori

thus joins the elite of stars with the best established absolute dimensions. The system is very young, and in fact it is so close to the ZAMS that within the observational errors the comparison with current stellar evolution models is unable to tell us whether it has already settled on the main sequence (age ~ 65 – 82 Myr, depending on the metallicity), or whether it is still at the end of the contraction phase (age = 7.7 Myr). We find good agreement with both scenarios for a chemical composition close to solar.

The tentative detection of apsidal motion in GG Ori with a period of $U = 10700$ yr has allowed us to make an initial test of interior structure models. This system is particularly interesting in that the contribution from General Relativity should be substantial ($\sim 70\%$ of the total apsidal motion). Although the agreement with theory is good, further observations to improve the error in $\dot{\omega}$ are necessary for a more stringent test.

Current mechanisms that describe the tidal evolution of binary properties disagree as to whether the rotation of the components of GG Ori should be synchronized with the orbital motion. Unfortunately our own measurements of $v \sin i$ have not settled the issue. Although this area of theory is perhaps the weakest of the comparisons we have discussed, the models do explain the significant eccentricity of the orbit system at the relatively short period of 6.6 days.

We thank P. Berlind, M. Calkins, D. W. Latham, A. Milone, and R. P. Stefanik, who obtained many of the spectroscopic observations used here, and R. Davis, who maintains the CfA database of radial velocities. We are also grateful to Daryl Willmarth at KPNO for assistance with the spectroscopic observations there. Paul Etzel provided helpful comments on aspects of the light curve solutions. We thank the referee for useful suggestions. This research has made use of the SIMBAD database, operated at CDS, Strasbourg, France, and of NASA’s Astrophysics Data System Bibliographic Services.

8. Appendix

The determination of the error in the relative radius of the secondary component, r_B , in eclipsing binaries has often been a source of confusion among users of EBOP. Results are occasionally published in which that uncertainty is derived by propagating errors in the expression $r_B = k \cdot r_A$, where k (defined as r_B/r_A), the primary radius r_A , and their associated uncertainties are usually adopted directly from the output of the program. Aside from the fact that the formal errors in the latter two quantities are typically underestimated (see §4), this procedure is incorrect, as already pointed out clearly in the documentation for EBOP (Etzel 1980). The reason is that k and r_A are, as a rule, strongly (negatively)

correlated, and simply adding their uncertainties by quadratures does not account for this correlation. It is still possible to make correct use of the expression above if the correlation is explicitly taken into account by adding the corresponding term from the off-diagonal elements of the covariance matrix in the least squares solution. However, this information is not readily available in the standard output from EBOP. Alternative procedures for estimating σ_{r_B} are discussed in the documentation for the code (see also Popper & Etzel 1981). For example, when k is fixed in the least squares solution one may use as an approximation the expression

$$\sigma_{r_B} = k \cdot \sigma_{r_A}, \quad (1)$$

although this explicitly ignores the error in k , which in our case is well known independently of EBOP.

The reason for the strong correlation between k and r_A is that the *sum* of the radii, $r \equiv r_A + r_B$, which is directly related to the form of the light curve, is usually very well determined, and therefore if k is increased the primary radius r_A must decrease to maintain the sum. This suggests that a more sensible way to approach the problem might be to consider the expressions

$$r_A = \left(\frac{1}{1+k} \right) r, \quad r_B = \left(\frac{k}{1+k} \right) r, \quad (2)$$

in which the independent variables are r and k , which are *not* strongly correlated in general. For GG Ori we have shown this to be the case in §4. Propagation of errors in these equations may therefore be expected to give more reasonable estimates of σ_{r_A} and σ_{r_B} :

$$\sigma_{r_A} = \frac{1}{1+k} \left[\left(\frac{r}{1+k} \right)^2 \sigma_k^2 + \sigma_r^2 \right]^{1/2}, \quad \sigma_{r_B} = \frac{1}{1+k} \left[\left(\frac{r}{1+k} \right)^2 \sigma_k^2 + k^2 \sigma_r^2 \right]^{1/2}. \quad (3)$$

Note that σ_{r_B} will be smaller or larger than σ_{r_A} depending only on the value of k , as in eq.(1). In fact, setting $\sigma_k = 0$ and combining the expressions above leads back to (1).

Practical application of these equations requires an estimate of σ_r . Although r is not one of the adjustable variables in EBOP, it is not difficult to obtain an estimate of its error by experimenting with the light curve solutions and assessing the sensitivity of r to the various input quantities, as we have in fact done with r_A . However, to ensure consistency with the error in r_A , we have chosen to use the first of the equations in (3) to solve for σ_r , since σ_{r_A} is already known and so is σ_k . Once we determined the value of σ_r needed to reproduce σ_{r_A} , we used it in the expression for σ_{r_B} .

REFERENCES

- Andersen, J. 1991, *A&A Rev.*, 3, 91
- Andersen, J. 1998, in *Fundamental Stellar Properties: The Interaction between Observation and Theory*, IAU Symp. 189, eds. T. R. Bedding, A. J. Booth & J. Davis (Dordrecht: Reidel), 99
- Blaauw, A. 1964, *ARA&A*, 2, 213
- Braune, W., Hübscher, J., & Mundry, E. 1977, *AN*, 298, 121
- Brown, A. G. A., de Geus, E. J., & de Zeeuw, P. T. 1994, *A&A*, 289, 101
- Brown, A. G. A., Walter, F. M., & Blaauw, A. 1999, in *The Orion Complex Revisited*, ASP Conf. Ser., ed. M. J. McCaughrean & A. Burkert (San Francisco: ASP), in press
- Busch, H. 1975, *Mitt. der Bruno-H.-Burgel Sternwarte Hartha*, 8, 18
- Casey, B. W., Mathieu, R. D., Vaz, L. P. R., Andersen, J., & Suntzeff, N. B. 1998, *AJ*, 115, 1617
- Claret, A. 1995, *A&AS*, 109, 441
- Claret, A., & Cunha, N. C. S. 1997, *A&A*, 318, 187
- Claret, A., & Giménez, A. 1992, *A&AS*, 96, 255
- Claret, A., Giménez, A., & Cunha, N. C. S. 1995, *A&A*, 299, 724
- Claret, A., Giménez, A., & Martín, E. L. 1995, *A&A*, 302, 741
- Clausen, J. V., & Nördstrom, B. 1980, *A&A*, 83, 339
- Crawford, D. L. 1978, *AJ*, 83, 48
- Deutschman, W. A., Davis, R. J., & Schild, R. E. 1976, *ApJS*, 30, 97
- Diethelm, R. 1975, *BBSAG Bull.*, 21, 4
- Diethelm, R. 1976, *BBSAG Bull.*, 25, 2
- Diethelm, R. 1983, *BBSAG Bull.*, 65, 4
- Diethelm, R. 1992, *BBSAG Bull.*, 21, 1
- Duquennoy, A., Mayor, M., & Mermilliod, J.-C. 1992, in *Binaries as Tracers of Stellar Formation*, eds. A. Duquennoy & M. Mayor (Cambridge: Cambridge Univ. Press), 52
- Etzel, P. B. 1980, *EBOP User's Guide*, Univ. of California, Los Angeles
- Etzel, P. B. 1981, in *Photometric and Spectroscopic Binary Systems*, ed. E. B. Carling & Z. Kopal (Dordrecht: Reidel), 111

- Fekel, F. C. 1999, in *Precise Radial Velocities*, ASP Conf. Ser. 185, eds. J. B. Hearnshaw & C. D. Scarfe (San Francisco: ASP), 378
- Giménez, A. 1985, *ApJ*, 297, 405
- Hilditch, R. W., & Hill, G. 1975, *MmRAS*, 79, 101
- Hoffleit, D. 1982, *The Bright Star Catalogue* (New Haven: Yale Univ. Obs.)
- Hoffmeister, C. 1934, *AN*, 253, 195
- Izotov, Y. I., & Thuan, T. X. 1998, *ApJ*, 500, 188
- Johnson, H. L., & Morgan, W. W. 1953, *ApJ*, 117, 313
- Kholopov, P. N. 1985, *General Catalog of Variable Stars* (Moscow: Nauka), Vol. II
- Kordylewski, K. 1951, *Acta Astr. Ser. C*, 4, 134
- Lacy, C. H. S. 1984, *IBVS*, No. 2489
- Lacy, C. H. S. 1992a, *AJ*, 104, 801
- Lacy, C. H. S. 1992b, *AJ*, 104, 2213
- Lacy, C. H. S., & Fox, G. W. 1994, *IBVS*, No. 4009
- Lacy, C. H. S., Ibanoglu, C., Tunca, Z., Evren, S., Akan, C., Keskin, V., Zakirov, M., Arzumanyants, G., Ishankulov, R., & Kharchenko, V. 1995, *IBVS*, No. 4194
- Lacy, C. H. S., Torres, G., Claret, A., Stefanik, R. P., Latham, D. W., & Sabby, J. A. 2000, *AJ*, 119, 1389
- Landolt, A. U. 1973, *AJ*, 78, 959
- Latham, D. W. 1992, in *IAU Coll. 135, Complementary Approaches to Double and Multiple Star Research*, ASP Conf. Ser. 32, eds. H. A. McAlister & W. I. Hartkopf (San Francisco: ASP), 110
- Latham, D. W., Nordström, B., Andersen, J., Torres, G., Stefanik, R. P., Thaller, M., & Bester, M. 1996, *A&A*, 314, 864
- Lavrov, M. I. 1993, *Trudy Kazansk. Gor. Astron. Obs.*, 53, 34
- Mamajek, E. E., Lawson, W. A., Feigelson, E. D. 2000, *ApJ*, in press
- Martynov, D. Y., 1973, in *Eclipsing Variable Stars*, ed. V. P. Tsesevich (New York: Halsted), 146
- Mathieu, R. D., Duquennoy, A., Latham, D. W., Mayor, M., Mazeh, T., & Mermilliod, J.-C. 1992, in *Binaries as Tracers of Stellar Formation*, ed. A. Duquennoy & M. Mayor (Cambridge: Cambridge Univ. Press), 278

- Nikulina, T. G. 1958, *Astr. Circ.*, 189, 16
- Pagel, B. E. J., & Portinari, L. 1998, *MNRAS*, 298, 747
- Peimbert, M. 1993, *Rev. Mex. Astr. Astrof.*, 27, 9
- Pols, O. R., Tout, C. A., Schröder, K.-P., Eggleton, P. P., & Manners, J. 1997, *MNRAS*, 289, 869
- Popper, D. M. 1968, *ApJ*, 154, 191
- Popper, D. M. 1987, *ApJ*, 313, L81
- Popper, D. M. 1980, *ARA&A*, 18, 115
- Popper, D. M., & Etzel, P. B. 1981, *AJ*, 86, 102
- Renzini, A. 1994, *A&A*, 285, L5
- Ribas, I., Jordi, C., Torra, J., & Giménez, A. 2000, *MNRAS*, 313, 99
- Soloviev, A. 1945, *Astr. Circ.*, 41, 8
- Stefanik, R. P., Latham, D. W., & Torres, G. 1999, in *Precise Stellar Radial Velocities*, IAU Coll. 170, ASP Conf. Ser., 185, eds. J. B. Hearnshaw & C. D. Scarfe (San Francisco: ASP), 354
- Szafraniec, R. 1952, *Acta Astr. Ser. C*, 5, 7
- Szafraniec, R. 1955, *Acta Astr. Ser. C*, 5, 189
- Szafraniec, R. 1956, *Acta Astr.*, 6, 141
- Szafraniec, R. 1958, *Acta Astr.*, 8, 189
- Tassoul, M., & Tassoul, J.-L. 1997, *ApJ*, 481, 363
- Tshuprina, P. I. 1957, *Astr. Circ.*, 183, 16
- Torres, G., Andersen, J., Nördstrom, B., & Latham, D. W. 2000, *AJ*, 119, 1942
- Torres, G., Lacy, C. H. S., Claret, A., Zakirov, M. M., Arzumanyants, G. C., Bayramov, N., Hojaev, A. S., Stefanik, R. P., Latham, D. W., & Sabby, J. A. 1999, *AJ*, 118, 1831
- Torres, G., Stefanik, R. P., Andersen, J., Nördstrom, B., Latham, D. W., & Clausen, J. V. 1997, *AJ*, 114, 2764
- Verbunt, F., & Phinney, E. S. 1995, *A&A*, 296, 709
- Winiarski, M. 1972, *Acta Astr.*, 24, 89
- Wade, R. A., & Rucinski, S. M. 1985, *A&AS*, 60, 471
- Warren, W. H. Jr., & Hesser, J. E. 1977, *ApJS*, 34, 115

- Zakirov, M. M. 1997, *Pis'ma Astr. Zh.*, 23, 626
- Zahn, J.-P. 1992, in *Binaries as Tracers of Stellar Formation*, ed. A. Duquennoy & M. Mayor (Cambridge: Cambridge Univ. Press), 253
- Zahn, J.-P., & Bouchet, L. 1989, *A&A*, 223, 112
- de Zeeuw, P. T., Hoogerwerf, R., de Bruijne, J. H. J., Brown, A. G. A., & Blaauw, A. 1999, *AJ*, 117, 354
- Zessewitsch, W. P. 1945, *Astr. Circ.*, 40, 7
- Zucker, S., & Mazeh, T. 1994, *ApJ*, 420, 806




Fig. 1.— Radial-velocity curves of GG Ori from our final spectroscopic solution that includes times of minimum in the least-squares fit. The filled circles are for the CfA observations of the primary, the open circles are for the secondary, and the asterisks represent the velocity measurements from KPNO.




Fig. 2.— a) Ephemeris curve for GG Ori along with all available times of eclipse (solid line for the primary); b) enlargement to show the distribution of the measurements. The time distribution of the spectroscopic observations included in the apsidal motion analysis is indicated by the horizontal line in the upper right.




Fig. 3.— Application of the spectroscopic constraints on the light ratio in B and V to determine k , the ratio of the radii of the components. The solid curves are derived from light curve solutions based on our CTIO photometry with k held fixed. The arrow with the error bar marks the resulting weighted average of k .




Fig. 4.— Light curves obtained at CTIO in the B and V bands along with the fitted solutions from EBOP. The lower panels expand the region near the primary and secondary minima for the V band.




Fig. 5.— Light curves in $UBVR$ based on the observations by Zakirov (1997), along with our fits.




Fig. 6.— Comparison with stellar evolution models for both components in the $\log g$ – $\log T_{\text{eff}}$ plane. Evolutionary tracks for two different compositions are shown, giving good fits assuming GG Ori is on the main sequence (MS). The uncertainty in the placement of the tracks due to errors in the measured masses is indicated with the error bar in the lower left. The full extent of the main sequence band is shown in the inset, emphasizing the fact that the system is very close to the ZAMS.




Fig. 7.— Best fit evolutionary tracks assuming GG Ori is still in the pre-main sequence (PMS) phase. The uncertainty in the placement of the tracks due to errors in the measured masses is indicated with the error bar in the lower right.




Fig. 8.— Evolutionary track for a mass equal to that of the primary in GG Ori and a composition close to solar ($Z = 0.020$, $Y = 0.2729$). Also shown are all other binary systems that have accurate absolute dimensions available in which at least one component has a similar mass. The 8 stars represented differ in mass by less than 1%, and all are adequately fit by this model.




Fig. 9.— Critical values of the surface gravity at which the stars in GG Ori reach synchronization and circularization, as a function of orbital period. Curves for two different tidal mechanisms are shown: Zahn (1992) and Tassoul & Tassoul (1997). Evolution carries the stars upwards. Our measurements of $\log g$ for the system are indicated with crosses. Both theories predict circularization has not yet occurred (as observed), but they disagree as to the synchronization.

TABLE 1. KPNO radial velocity measurements of GG Ori, and residuals from the final spectroscopic orbit.

HJD (2,400,000+)	RV _A (km s ⁻¹)	RV _B (km s ⁻¹)	(O-C) _A (km s ⁻¹)	(O-C) _B (km s ⁻¹)	Orbital phase
45684.8406.....	+91.5	-56.9	+3.99	+0.53	0.394
46070.8049.....	-60.8	+89.1	-0.99	-2.37	0.087
46071.8576.....	-79.4	+108.3	+2.60	-0.21	0.398
47778.9875.....	+82.3	-54.6	-1.12	-3.71	0.906
47779.9970.....	+98.3	-70.3	-0.41	-0.05	0.199
47781.9962.....	-85.1	+110.9	-1.45	-0.83	0.507
48150.9916.....	+96.3	-72.1	-2.42	+1.86	0.236
48152.0011.....	+71.8	-43.1	-0.79	-2.71	0.027
51243.7258.....	-87.7	+118.1	+0.96	+2.88	0.167
51246.7258.....	+60.8	-33.1	-0.36	+2.02	0.168
51247.6287.....	+94.6	-65.4	+0.47	+0.03	0.162

TABLE 2. CfA radial velocity measurements of GG Ori, and residuals from the final spectroscopic orbit.

HJD (2,400,000+)	RV _A ^a (km s ⁻¹)	RV _B ^a (km s ⁻¹)	(O-C) _A (km s ⁻¹)	(O-C) _B (km s ⁻¹)	Orbital phase
50144.6930.....	-9.01	+39.00	+0.79	+0.53	0.394
50407.9143.....	-73.46	+97.71	-2.16	-2.37	0.087
50409.9760.....	-10.06	+36.68	-1.84	-0.21	0.398
50439.8695.....	+69.42	-47.59	-2.98	-3.71	0.906
50441.8148.....	-90.88	+119.08	-0.57	-0.05	0.199
50443.8593.....	+35.21	-9.50	-2.04	-0.83	0.507
50448.6911.....	-79.78	+110.30	-0.13	+1.86	0.236
50473.8339.....	-22.15	+48.85	+0.71	-2.71	0.027
50474.7629.....	-93.01	+126.09	+1.37	+2.88	0.167
50494.6588.....	-94.59	+125.22	-0.21	+2.02	0.168
50739.9868.....	-93.01	+123.32	+1.46	+0.03	0.162
50770.8719.....	+97.91	-66.73	+0.27	+2.43	0.819
50772.9141.....	-89.25	+118.60	+0.44	+0.10	0.127
50792.9273.....	-95.67	+117.64	-2.32	-4.53	0.145
50799.8040.....	-93.11	+125.32	+0.10	+3.29	0.182
50799.8119.....	-91.45	+122.44	+1.60	+0.56	0.183
50823.8998.....	+95.85	-68.85	-2.18	+0.71	0.816
50825.8988.....	-88.51	+117.77	-2.11	+2.56	0.117
50826.7972.....	-73.96	+103.43	-0.53	+1.22	0.253
50826.8636.....	-68.20	+99.13	+1.21	+0.94	0.263
50826.8738.....	-68.02	+100.13	+0.75	+2.58	0.264
50859.7244.....	-86.99	+113.14	-1.50	-1.16	0.218
50916.6656.....	+99.47	-73.36	+0.48	-2.85	0.804
51089.0199.....	+100.83	-72.47	+1.35	-1.46	0.795
51117.9532.....	-95.72	+124.68	-1.33	+1.46	0.158
51121.9141.....	+100.86	-68.75	+2.02	+1.62	0.755
51148.8862.....	+98.99	-69.36	+1.71	-0.56	0.822
51177.7818.....	-91.89	+124.19	+1.64	+1.84	0.180
51241.6644.....	+100.05	-71.21	+1.74	-1.38	0.813
51267.6205.....	+96.22	-68.22	+0.04	-0.52	0.727
51268.6168.....	+87.27	-55.87	+2.77	+0.13	0.877
51292.6444.....	+36.14	-5.90	+1.51	+0.14	0.500
51472.9868.....	+88.29	-61.96	-3.13	+0.97	0.695
51474.0126.....	+89.66	-61.47	-2.73	+2.44	0.850
51475.9391.....	-94.76	+123.06	-2.13	+1.60	0.140
51533.8139.....	+88.15	-58.59	+0.54	+0.53	0.868
51539.8100.....	+101.68	-69.24	+2.09	+1.88	0.772
51566.8594.....	+93.02	-62.84	+0.84	+0.86	0.851
51569.7008.....	-63.96	+92.21	-1.54	+1.03	0.279
51592.8330.....	+93.60	-69.04	-5.86	+1.95	0.767
51619.6948.....	+95.01	-68.30	-2.74	+0.98	0.818
51621.7149.....	-89.20	+114.03	-0.89	-3.10	0.123

^aIncludes corrections described in the text.

TABLE 3. Spectroscopic orbital solutions for GG Ori.

Element (1)	KPNO (2)	CfA (3)	Combined (4)
Adjusted quantities			
P_s (days) ^a	6.6314948	6.6314948	6.6314936 ± 0.0000017
γ (km s ⁻¹).....	$+14.13 \pm 0.35$	$+14.30 \pm 0.21$	$+14.31 \pm 0.21$
ΔRV (km s ⁻¹) ^b	...	...	-0.06 ± 0.39
K_A (km s ⁻¹).....	96.80 ± 0.71	97.17 ± 0.34	97.10 ± 0.30
K_B (km s ⁻¹).....	96.67 ± 0.51	97.53 ± 0.33	97.28 ± 0.28
e	0.2264 ± 0.0039	0.2201 ± 0.0030	0.2218 ± 0.0022
ω_A (deg).....	122.7 ± 1.0	123.34 ± 0.67	122.76 ± 0.39
$\dot{\omega}$ (deg cycle ⁻¹).....	...	...	0.00061 ± 0.00025
Min I (HJD–2,400,000) ^a ..	49,717.66253	49,717.66253	$49,717.66259 \pm 0.00020$
Derived quantities			
P_a (days).....	...	...	6.6315048 ± 0.0000042
U (yr).....	...	...	10700 ± 4500
T (HJD–2,400,000) ^c	$49,717.999 \pm 0.025$	$49,718.073 \pm 0.012$	$49,718.0448 \pm 0.0078$
$a_A \sin i$ (10 ⁶ km).....	8.597 ± 0.073	8.644 ± 0.032	8.634 ± 0.028
$a_B \sin i$ (10 ⁶ km).....	8.586 ± 0.054	8.676 ± 0.031	8.650 ± 0.026
$a \sin i$ (R _☉).....	24.69 ± 0.13	24.885 ± 0.065	24.833 ± 0.056
$M_A \sin^3 i$ (M _☉).....	2.297 ± 0.036	2.358 ± 0.019	2.341 ± 0.016
$M_B \sin^3 i$ (M _☉).....	2.300 ± 0.043	2.349 ± 0.020	2.337 ± 0.017
$q \equiv M_B/M_A$	1.001 ± 0.010	0.9963 ± 0.0051	0.9982 ± 0.0044
Other quantities pertaining to the fit			
N_{obs}	11	42	53
$N_{\text{min I}}$	...	...	24
$N_{\text{min II}}$	...	...	32
Time span (yr) ^d	15.2	4.0	70.1
σ_A (km s ⁻¹).....	1.93	1.95	1.83/1.93
σ_B (km s ⁻¹).....	1.38	1.91	1.33/1.94

^aSidereal period and epoch fixed from §3 for the KPNO and CfA solutions, and adjusted for the combined solution.

^bVelocity offset in the sense ⟨CfA–KPNO⟩.

^cPeriastron passage closest to the epoch of primary minimum.

^dIncludes times of minimum for the combined solution.

TABLE 4. Eclipse timings and residuals for GG Ori.

HJD (2,400,000+)	Year	Eclipse ^a	Type ^b	Epoch ^c	(O-C) (days)	Reference
26003.4590	1930.073	1	pg	−3576.0	+0.0312	1
26384.2800	1931.115	2	pg	−3519.0	+0.0258	1
27100.4170	1933.076	2	pg	−3411.0	−0.0380	1
27787.3050	1934.956	1	v	−3307.0	+0.0044	2
27823.3050	1935.055	2	v	−3302.0	+0.0177	2
28165.2900	1935.991	1	v	−3250.0	−0.0059	2
28181.3900	1936.035	2	v	−3248.0	+0.0023	2
29697.1430	1940.185	1	v	−3019.0	−0.0288	3
29697.1440	1940.185	1	pg	−3019.0	−0.0278	4
30022.1290	1941.075	1	v	−2970.0	+0.0138	3
30022.1330	1941.075	1	pg	−2970.0	+0.0178	4
31156.1200	1944.180	1	v	−2799.0	+0.0187	5
31477.2900	1945.059	2	v	−2751.0	+0.0523	5
32969.3400	1949.144	2	v	−2526.0	+0.0173	2
32973.1540	1949.154	1	pg	−2525.0	+0.0224	4
33006.2900	1949.245	1	v	−2520.0	+0.0009	2
33181.4720	1949.725	2	pg	−2494.0	−0.0584	4
33274.3420	1949.979	2	pg	−2480.0	−0.0292	4
33307.5300	1950.070	2	v	−2475.0	+0.0014	2
33311.3300	1950.080	1	v	−2474.0	−0.0080	2
33334.1340	1950.143	2	pg	−2471.0	+0.0794	4
33596.4910	1950.861	1	v	−2431.0	−0.0013	6
33689.3380	1951.115	1	v	−2417.0	+0.0047	6
34034.1310	1952.059	1	pg	−2365.0	−0.0402	4
34448.1660	1953.193	2	pg	−2303.0	+0.0213	4
35071.5100	1954.899	2	v	−2209.0	+0.0053	7
35164.3380	1955.154	2	pg	−2195.0	−0.0076	8
35376.5260	1955.734	2	v	−2163.0	−0.0272	9
35429.6040	1955.880	2	pg	−2155.0	−0.0011	8
35509.1850	1956.098	2	pg	−2143.0	+0.0020	4
35542.3360	1956.188	2	pg	−2138.0	−0.0044	8
35807.5590	1956.915	2	v	−2098.0	−0.0410	10
35867.2770	1957.078	2	pg	−2089.0	−0.0064	8
35904.2090	1957.179	1	pg	−2083.0	−0.0445	4
35904.2600	1957.179	1	v	−2083.0	+0.0065	10
38327.5220	1963.814	2	pg	−1718.0	−0.0438	11
38384.4720	1963.970	1	pg	−1709.0	+0.0385	11
38440.3050	1964.123	2	pg	−1701.0	+0.0039	11
38457.3380	1964.169	1	pg	−1698.0	−0.0420	11
38652.5880	1964.704	2	pg	−1669.0	+0.0792	11
41391.3470	1972.202	2	pg	−1256.0	+0.0333	11
41676.4660	1972.983	2	pg	−1213.0	−0.0017	12
41961.5670	1973.763	2	pg	−1170.0	−0.0548	11
42074.3590	1974.072	2	pg	−1153.0	+0.0019	11
42452.3650	1975.107	2	pg	−1096.0	+0.0130	11
42469.2960	1975.153	1	v	−1093.0	(−0.1399)	13
42469.3720	1975.154	1	v	−1093.0	−0.0639	14
42777.3060	1975.997	2	v	−1047.0	+0.0111	15
45403.3370	1983.186	2	v	−651.0	−0.0276	16
48590.3086	1991.912	1	pe	−170.0	+0.0006	17
48606.3690	1991.956	2	v	−168.0	−0.0048	18
48623.4634	1992.003	1	v	−165.0	−0.0021	18
48911.4212	1992.791	2	pe	−122.0	−0.0011	17
48948.4090	1992.892	1	pe	−116.0	+0.0001	17
49355.7327	1994.007	2	pe	−55.0	+0.0006	19
49366.1968	1994.036	1	pe	−53.0	+0.0036	17
49717.6624	1994.998	1	pe	0.0	−0.0002	17

^a1 = primary eclipse; 2 = secondary eclipse.

^bpg = photographic, v = visual, pe = photoelectric.

^cCounted from primary minimum in column 4 of Table 3.

References for Table 4.

- (1) Hoffmeister 1934; (2) Kordylewski 1951; (3) Soloviev 1945; (4) Nikulina 1958; (5) Zesewitsch 1945; (6) Szafraniec 1952; (7) Szafraniec 1955; (8) Tshuprina 1957; (9) Szafraniec 1956; (10) Szafraniec 1958; (11) Busch 1975; (12) Winiarski 1972; (13) Braune et al. 1977; (14) Diethelm 1975; (15) Diethelm 1976; (16) Diethelm 1983; (17) Lacy et al. 1995; (18) Diethelm 1992; (19) Lacy & Fox 1994.

TABLE 5. Absolute photometry of GG Ori and comparison stars.

Star	V	$B - V$	$U - B$	N_{obs}
GG Ori (1988-1990)	10.359 ± 0.009	0.532 ± 0.004	0.323 ± 0.005	4
GG Ori (1993-1995)	10.385 ± 0.007	0.490 ± 0.003	0.326 ± 0.005	3
BD-1°1013	8.802 ± 0.003	0.232 ± 0.001	-0.224 ± 0.004	2
BD-0°1081	8.960 ± 0.004	0.350 ± 0.004	0.027 ± 0.002	3
BD-0°1044	7.341	0.523	-0.026	1

TABLE 6. Differential photometric observations of GG Ori from CTIO in the B band, in the sense $\langle \text{variable} - \text{comparison} \rangle$.

HJD (2,400,000+)	B (mag)	HJD (2,400,000+)	B (mag)	HJD (2,400,000+)	B (mag)	HJD (2,400,000+)	B (mag)
49339.6586	2.536	49342.5826	2.038	49355.5552	1.884	49717.6308	2.333
49339.6677	2.567	49342.5859	2.036	49355.5594	1.897	49717.6353	2.382
49339.6806	2.497	49343.5543	1.867	49355.5625	1.903	49717.6391	2.408
49339.6869	2.444	49343.5590	1.886	49355.5658	1.900	49717.6427	2.439
49339.6919	2.405	49343.5637	1.889	49355.5909	1.956	49717.6462	2.474
49339.6966	2.360	49344.6894	1.859	49355.5953	1.938	49717.6498	2.492
49339.7145	2.221	49344.6961	1.876	49355.5986	1.966	49717.6533	2.524
49339.7190	2.184	49344.7014	1.866	49355.6020	1.974	49717.6569	2.545
49339.7239	2.162	49348.6986	1.870	49355.6052	1.991	49717.6613	2.543
49339.7283	2.116	49348.7052	1.871	49355.6274	2.058	49717.6648	2.554
49339.7339	2.099	49348.7102	1.873	49355.6318	2.059	49717.6684	2.543
49339.7385	2.067	49348.8059	1.843	49355.6353	2.088	49717.6721	2.527
49339.7446	2.023	49348.8101	1.825	49355.6388	2.110	49717.6757	2.486
49339.7487	2.010	49349.5506	1.931	49355.6438	2.127	49717.6793	2.464
49339.7527	1.990	49349.5606	1.887	49355.6487	2.140	49717.6840	2.432
49339.7607	1.941	49349.5673	1.868	49355.6730	2.260	49717.6872	2.402
49339.7684	1.923	49349.6409	1.881	49355.6775	2.274	49717.6907	2.371
49339.7754	1.894	49349.6454	1.889	49355.6807	2.304	49717.6943	2.341
49339.7820	1.882	49349.6488	1.886	49355.6840	2.316	49717.6978	2.296
49339.7886	1.868	49349.6533	1.882	49355.6874	2.343	49717.7037	2.262
49339.7944	1.858	49349.6566	1.889	49355.6907	2.357	49717.7073	2.223
49339.8041	1.852	49349.6610	1.878	49355.6950	2.389	49717.7111	2.223
49339.8086	1.855	49349.6642	1.876	49355.6983	2.386	49717.7146	2.179
49339.8138	1.854	49349.6683	1.861	49355.7017	2.419	49717.7192	2.149
49339.8181	1.854	49349.6716	1.866	49355.7050	2.417	49717.7228	2.136
49339.8225	1.856	49349.7644	1.862	49355.7086	2.454	49717.7264	2.104
49339.8267	1.837	49349.7688	1.862	49355.7132	2.467	49717.7304	2.071
49339.8331	1.881	49349.7722	1.864	49355.7166	2.465	49717.7340	2.059
49340.5368	1.829	49349.7755	1.863	49355.7201	2.493	49717.7376	2.028
49340.5453	1.926	49349.7812	1.857	49355.7234	2.505	49717.7412	2.019
49340.5520	1.908	49349.7847	1.856	49355.7270	2.509	49717.7447	1.992
49340.5578	1.919	49350.5867	1.897	49355.7305	2.506	49717.7485	1.973
49340.5639	1.896	49350.5902	1.879	49355.7341	2.514	49717.7563	1.940
49340.5680	1.884	49350.5933	1.892	49355.7376	2.502	49717.7626	1.915
49340.5736	1.889	49350.6623	1.891	49355.7421	2.502	49717.7681	1.899
49340.5776	1.884	49350.6686	1.894	49355.7455	2.488	49717.7750	1.866
49340.5821	1.894	49350.6762	1.880	49355.7494	2.477	49717.7799	1.863
49340.5862	1.889	49350.7574	1.861	49355.7536	2.453	49717.7862	1.859
49340.5915	1.897	49350.7620	1.855	49355.7571	2.436	49717.7972	1.849
49340.5955	1.878	49350.7654	1.859	49716.5883	1.878	49720.5503	2.058
49340.6794	1.870	49350.8027	1.861	49716.5977	1.884	49720.5547	2.154
49340.6857	1.869	49350.8074	1.838	49716.6064	1.885	49720.5556	2.138
49340.6898	1.869	49350.8114	1.837	49716.7719	1.873	49720.5601	2.130
49340.6943	1.868	49352.7154	1.867	49716.7784	1.847	49720.5646	2.100
49340.6982	1.878	49352.7214	1.866	49716.7831	1.850	49720.5680	2.097
49340.7018	1.878	49352.7266	1.858	49717.5705	1.971	49720.5713	2.053
49340.7573	1.868	49352.7598	1.846	49717.5757	1.991	49720.5755	2.047
49340.7661	1.857	49352.7644	1.861	49717.5799	2.010	49720.5788	2.036
49340.7726	1.866	49352.7684	1.860	49717.5846	2.029	49720.5823	2.017
49341.7584	1.873	49352.7721	1.859	49717.5883	2.048	49720.5859	2.000
49341.7664	1.860	49353.5467	1.883	49717.5920	2.066	49720.5891	2.000
49341.7731	1.860	49353.5515	1.899	49717.5958	2.085	49720.5934	1.984
49342.5482	2.194	49353.5549	1.890	49717.6005	2.127	49720.5969	1.967
49342.5544	2.155	49353.5585	1.892	49717.6041	2.151	49720.6021	1.956
49342.5590	2.158	49354.6393	1.874	49717.6080	2.160	49720.6071	1.949
49342.5635	2.116	49354.6439	1.869	49717.6116	2.192	49720.7236	1.880
49342.5668	2.114	49354.6474	1.868	49717.6164	2.243	49720.7289	1.874
49342.5714	2.096	49355.5410	1.891	49717.6198	2.253	49720.7326	1.867
49342.5747	2.071	49355.5468	1.884	49717.6233	2.280	49720.7814	1.865
49342.5791	2.062	49355.5521	1.890	49717.6271	2.320	49720.7873	1.858
49720.7913	1.854	49722.6534	1.882	49722.7456	1.855	49727.5643	1.906
49721.7524	1.849	49722.6587	1.870	49726.6865	1.878	49727.5700	1.908
49721.7590	1.857	49722.7343	1.874	49726.6927	1.891		
49721.7643	1.849	49722.7378	1.854	49726.6981	1.883		
49722.6461	1.883	49722.7423	1.852	49727.5591	1.907		

TABLE 7. Differential photometric observations of GG Ori from CTIO in the V band, in the sense $\langle \text{variable} - \text{comparison} \rangle$.

HJD (2,400,000+)	V (mag)	HJD (2,400,000+)	V (mag)	HJD (2,400,000+)	V (mag)	HJD (2,400,000+)	V (mag)
49339.6590	2.236	49342.5830	1.754	49355.5555	1.605	49717.6312	2.074
49339.6681	2.291	49342.5863	1.739	49355.5597	1.607	49717.6357	2.096
49339.6810	2.224	49343.5546	1.579	49355.5629	1.626	49717.6395	2.144
49339.6873	2.164	49343.5594	1.606	49355.5662	1.620	49717.6430	2.161
49339.6923	2.124	49343.5641	1.609	49355.5913	1.672	49717.6465	2.197
49339.6970	2.088	49344.6898	1.604	49355.5957	1.683	49717.6502	2.220
49339.7149	1.947	49344.6965	1.593	49355.5990	1.688	49717.6537	2.242
49339.7194	1.916	49344.7018	1.600	49355.6024	1.691	49717.6573	2.268
49339.7243	1.878	49348.6990	1.601	49355.6055	1.709	49717.6617	2.277
49339.7287	1.856	49348.7056	1.589	49355.6278	1.790	49717.6652	2.277
49339.7343	1.823	49348.7106	1.585	49355.6322	1.808	49717.6689	2.267
49339.7389	1.792	49348.8063	1.586	49355.6357	1.813	49717.6725	2.235
49339.7450	1.754	49348.8105	1.572	49355.6392	1.826	49717.6761	2.211
49339.7491	1.741	49349.5509	1.632	49355.6442	1.841	49717.6797	2.181
49339.7531	1.709	49349.5610	1.602	49355.6491	1.874	49717.6844	2.143
49339.7611	1.679	49349.5677	1.601	49355.6734	1.986	49717.6876	2.124
49339.7688	1.648	49349.6413	1.591	49355.6779	2.016	49717.6911	2.088
49339.7758	1.626	49349.6458	1.596	49355.6811	2.025	49717.6947	2.059
49339.7824	1.601	49349.6492	1.609	49355.6844	2.042	49717.6982	2.027
49339.7890	1.595	49349.6537	1.600	49355.6878	2.061	49717.7041	1.983
49339.7948	1.599	49349.6570	1.605	49355.6911	2.074	49717.7077	1.957
49339.8045	1.591	49349.6614	1.585	49355.6954	2.097	49717.7115	1.937
49339.8090	1.594	49349.6646	1.596	49355.6987	2.122	49717.7150	1.913
49339.8142	1.596	49349.6687	1.591	49355.7020	2.137	49717.7196	1.881
49339.8185	1.594	49349.6720	1.595	49355.7054	2.164	49717.7232	1.855
49339.8229	1.582	49349.7648	1.598	49355.7090	2.175	49717.7268	1.832
49339.8271	1.590	49349.7692	1.586	49355.7136	2.193	49717.7308	1.805
49339.8335	1.650	49349.7726	1.589	49355.7170	2.219	49717.7344	1.776
49340.5372	1.577	49349.7759	1.580	49355.7205	2.221	49717.7380	1.766
49340.5456	1.621	49349.7816	1.583	49355.7237	2.243	49717.7416	1.736
49340.5523	1.609	49349.7850	1.582	49355.7274	2.238	49717.7451	1.724
49340.5582	1.610	49350.5871	1.604	49355.7309	2.237	49717.7489	1.707
49340.5643	1.618	49350.5906	1.602	49355.7345	2.245	49717.7567	1.680
49340.5684	1.611	49350.5937	1.614	49355.7380	2.236	49717.7630	1.651
49340.5739	1.611	49350.6627	1.578	49355.7425	2.232	49717.7685	1.627
49340.5780	1.607	49350.6690	1.627	49355.7459	2.229	49717.7754	1.607
49340.5825	1.611	49350.6766	1.600	49355.7498	2.213	49717.7803	1.591
49340.5866	1.605	49350.7578	1.603	49355.7539	2.194	49717.7866	1.595
49340.5919	1.595	49350.7624	1.586	49355.7574	2.174	49717.7976	1.585
49340.5959	1.606	49350.7658	1.600	49716.5887	1.602	49720.5505	1.819
49340.6798	1.600	49350.8031	1.579	49716.5981	1.598	49720.5548	1.844
49340.6861	1.596	49350.8078	1.559	49716.6068	1.597	49720.5560	1.860
49340.6902	1.595	49350.8118	1.586	49716.7723	1.597	49720.5606	1.825
49340.6947	1.592	49352.7158	1.590	49716.7788	1.590	49720.5650	1.814
49340.6986	1.599	49352.7218	1.599	49716.7835	1.585	49720.5684	1.791
49340.7022	1.600	49352.7269	1.587	49717.5709	1.691	49720.5717	1.780
49340.7577	1.598	49352.7602	1.595	49717.5761	1.716	49720.5759	1.771
49340.7665	1.590	49352.7648	1.595	49717.5802	1.747	49720.5792	1.756
49340.7730	1.596	49352.7689	1.591	49717.5850	1.753	49720.5827	1.753
49341.7588	1.593	49352.7725	1.585	49717.5887	1.778	49720.5863	1.734
49341.7668	1.591	49353.5471	1.604	49717.5924	1.802	49720.5895	1.717
49341.7735	1.591	49353.5519	1.608	49717.5962	1.815	49720.5938	1.706
49342.5486	1.917	49353.5554	1.612	49717.6009	1.849	49720.5973	1.698
49342.5548	1.875	49353.5589	1.606	49717.6045	1.865	49720.6025	1.677
49342.5594	1.859	49354.6397	1.602	49717.6084	1.897	49720.6075	1.661
49342.5638	1.829	49354.6443	1.594	49717.6120	1.925	49720.7240	1.609
49342.5672	1.838	49354.6478	1.602	49717.6168	1.960	49720.7293	1.599
49342.5718	1.802	49355.5414	1.610	49717.6201	1.983	49720.7329	1.588
49342.5751	1.787	49355.5472	1.602	49717.6237	2.003	49720.7818	1.591
49342.5795	1.774	49355.5525	1.604	49717.6275	2.038	49720.7877	1.590
49720.7917	1.578	49722.6538	1.595	49722.7460	1.584	49727.5647	1.624
49721.7528	1.591	49722.6591	1.593	49726.6869	1.602	49727.5704	1.616
49721.7594	1.595	49722.7347	1.594	49726.6931	1.605		
49721.7647	1.592	49722.7382	1.583	49726.6985	1.609		
49722.6465	1.592	49722.7427	1.585	49727.5594	1.622		

TABLE 8. Light curve solutions for GG Ori based on our CTIO observations.

Parameter (1)	General Fits		Adopted Fits	
	B (2)	V (3)	B (4)	V (5)
J_B	0.991 ± 0.030	0.992 ± 0.018	1.002 ± 0.008	1.002 ± 0.005
r_A	0.074 ± 0.003	0.073 ± 0.002	0.0746 ± 0.0003	
$k \equiv r_B/r_A^{\text{a}}$	1.02 ± 0.08	1.04 ± 0.05	0.988	
i (deg)	89.20 ± 0.14	89.31 ± 0.11	89.24 ± 0.08	
$e \cos \omega$	-0.1187 ± 0.0003	-0.1193 ± 0.0002	-0.1191 ± 0.0003	
$e \sin \omega$	$+0.210 \pm 0.009$	$+0.198 \pm 0.005$	$+0.200 \pm 0.007$	
$u = u_A = u_B$	0.59 ± 0.12	0.48 ± 0.08	0.55 ± 0.05	0.55 ± 0.03
L_A	0.492 ± 0.043	0.482 ± 0.028	0.504 ± 0.006	0.504 ± 0.005
σ (mag)	0.01414	0.00879	0.01408	0.00884
N_{obs}	256	255	256	255

^aIn the final fits k was held fixed to the value derived from spectroscopic constraints described in the text.

TABLE 9. Light curve solutions based on the *UBVR* observations by Zakirov (1997).

Parameter	<i>U</i>	<i>B</i>	<i>V</i>	<i>R</i>
Restricted fits				
J_B	0.98 ± 0.04	1.006 ± 0.012	0.991 ± 0.008	1.006 ± 0.012
r_A	0.076 ± 0.002	0.0734 ± 0.0007	0.0746 ± 0.0005	0.0754 ± 0.0007
$k \equiv r_B/r_A^a$	0.988	0.988	0.988	0.988
i (deg)	89.5 ± 0.3	89.43 ± 0.09	89.35 ± 0.07	89.24 ± 0.09
$e \cos \omega$	-0.1197 ± 0.0013	-0.1202 ± 0.0002	-0.1197 ± 0.0002	-0.1194 ± 0.0002
$e \sin \omega$	$+0.180 \pm 0.024$	$+0.179 \pm 0.008$	$+0.178 \pm 0.006$	$+0.196 \pm 0.008$
$u = u_A = u_B$	0.49 (fixed)	0.36 ± 0.11	0.41 ± 0.09	0.33 ± 0.13
L_A	0.510 ± 0.034	0.503 ± 0.011	0.507 ± 0.008	0.503 ± 0.011
σ (mag)	0.02499	0.01837	0.01365	0.01774
N_{obs}	175	240	242	238
Adopted fits				
J_B	0.964 ± 0.016	1.004 ± 0.009	0.990 ± 0.007	1.009 ± 0.009
r_A		0.0746 ± 0.0005		
$k \equiv r_B/r_A^a$		0.988		
i (deg)		89.35 ± 0.07		
$e \cos \omega$		-0.1198 ± 0.0004		
$e \sin \omega$		$+0.184 \pm 0.008$		
$u = u_A = u_B$	0.49 (fixed)	0.41 ± 0.07	0.41 ± 0.05	0.23 ± 0.07
L_A	0.513 ± 0.011	0.504 ± 0.008	0.507 ± 0.008	0.502 ± 0.008
σ (mag)	0.02519	0.01848	0.01361	0.01781
N_{obs}	175	240	242	238

^aHeld fixed in all solutions, based on the spectroscopic constraints described in the text.

TABLE 10. Weighted average light elements for GG Ori.

Parameter	U	B	V	R
J_B	0.964 ± 0.016	1.003 ± 0.006	0.998 ± 0.004	1.009 ± 0.009
r_A		0.0746 ± 0.0010		
r_B^{a}		0.0737 ± 0.0010		
$r \equiv r_A + r_B^{\text{a}}$		0.1483 ± 0.0019		
$k \equiv r_B / r_A$		0.988 ± 0.009		
i (deg)		89.30 ± 0.10		
$e \cos \omega$		-0.1194 ± 0.0002		
$e \sin \omega$		$+0.193 \pm 0.005$		
$u = u_A = u_B$	0.49 (fixed)	0.50 ± 0.04	0.51 ± 0.03	0.23 ± 0.07
L_A	0.513 ± 0.011	0.504 ± 0.005	0.505 ± 0.004	0.502 ± 0.008

^aSee text for error estimates.

TABLE 11. Physical parameters of GG Ori.

Parameter	Primary	Secondary
Mass ($M_{\odot}$)	2.342 ± 0.016	2.338 ± 0.017
Radius ($R_{\odot}$)	1.852 ± 0.025	1.830 ± 0.025
$\log g$	4.272 ± 0.012	4.282 ± 0.012
$\bar{\rho}$ (gr cm^{-3})	0.519 ± 0.021	0.537 ± 0.022
$v \sin i$ (km s^{-1}) (KPNO)	24 ± 2	23 ± 2
$v \sin i$ (km s^{-1}) (CfA)	16 ± 1	16 ± 1
v_{sync} (km s^{-1}) (circular)	14.1 ± 0.2	14.0 ± 0.2
v_{sync} (km s^{-1}) (eccentric) ...	22.8 ± 0.3	22.5 ± 0.3
a ($R_{\odot}$)	24.835 ± 0.056	
$\log T_{\text{eff}}$	3.9978 ± 0.0088	3.9978 ± 0.0088
$\log L/L_{\odot}$	1.480 ± 0.037	1.470 ± 0.037
M_{bol} (mag)	1.020 ± 0.093	1.046 ± 0.092
F'_V	3.974 ± 0.003	3.974 ± 0.003
M_V (mag)	1.176 ± 0.038	1.202 ± 0.039
L_B/L_A	0.98 ± 0.11	
$m - M$ (mag)	8.20 ± 0.10	
Dist (pc)	438 ± 20	

TABLE 12. Evolutionary age of the components of GG Ori in the main-sequence (MS) and pre-main sequence (PMS) scenarios, in Myr.

Chemical composition	Primary	Secondary	Average
$Z = 0.020, Y = 0.2729$ (MS).....	93 ± 27	72 ± 27	82 ± 19
$Z = 0.021, Y = 0.2810$ (MS).....	75 ± 29	52 ± 32	65 ± 21
$Z = 0.022, Y = 0.2860$ (PMS)....	7.26	8.23	7.7